

# NOROVIRUS: INFECTION CONTROL AND MANAGEMENT - CHW

## POLICY<sup>®</sup>

### DOCUMENT SUMMARY/KEY POINTS

- The incubation period for *Norovirus*-associated gastroenteritis in humans is usually 24 to 48 hours, but cases can occur within 12 hours of exposure<sup>2</sup>.
- Symptoms usually last 24 to 60 hours.
- Stringent [contact and droplet precautions](#) are an effective way to terminate the transmission of the disease.
- If the child is vomiting or has significant diarrhoea, infectious droplets may facilitate viral spread. Therefore staff involved in direct care of a vomiting patient or one with diarrhoea should wear personal protective equipment including surgical masks for protection<sup>1</sup>. Ensure lid of toilet is closed when flushing.
- Hand hygiene with antiseptic wash and water is important to stop the spread to the healthcare worker, other patients and visitors.
- Patients with *Norovirus* should remain in the ward most appropriate to their medical condition where they can be best cared for. However, they **MUST** be nursed in a single room or cohorted with other children with *Norovirus* in a dedicated room with en-suite toilet and bathroom facilities.
- Cleaning of rooms, contaminated surfaces and toys needs to be done with hypochlorite disinfectant (dichloroisocyanurate tablets).

This document reflects what is currently regarded as safe practice. However, as in any clinical situation, there may be factors which cannot be covered by a single set of guidelines. This document does not replace the need for the application of clinical judgement to each individual presentation.

|                        |  |  |
|------------------------|--|--|
| <b>Approved by:</b>    | SCHN Policy, Procedure and Guideline Committee |  |
| <b>Date Effective:</b> | 1 <sup>st</sup> December 2014                  | <b>Review Period:</b> 3 years            |
| <b>Team Leader:</b>    | Clinical Nurse Consultant                      | <b>Area/Dept:</b> Infection Control, CHW |

## CHANGE SUMMARY

- Due for mandatory review.
- Added the following sections:
  - Hospital Volunteers
  - Linen
  - Outbreak Management

## READ ACKNOWLEDGEMENT

- Medical and Nursing staff caring for patients with or suspected of having Norovirus should read this document.
- Infection Control staff should read and acknowledge they understand the contents of this document.
- Pathology staff with contact with patients with or suspected of having Norovirus should read this document.

This document reflects what is currently regarded as safe practice. However, as in any clinical situation, there may be factors which cannot be covered by a single set of guidelines. This document does not replace the need for the application of clinical judgement to each individual presentation.

|                        |  |  |
|------------------------|--|--|
| <b>Approved by:</b>    | SCHN Policy, Procedure and Guideline Committee |  |
| <b>Date Effective:</b> | 1 <sup>st</sup> December 2014                  | <b>Review Period:</b> 3 years            |
| <b>Team Leader:</b>    | Clinical Nurse Consultant                      | <b>Area/Dept:</b> Infection Control, CHW |

# TABLE OF CONTENTS

|   |           |
|---|-----------|
| <b>Introduction</b> .....   | <b>4</b>  |
| <b>What is <i>Norovirus</i>?</b> .....  | <b>4</b>  |
| <b>Clinical Presentation</b> .....  | <b>4</b>  |
| <b>Command and Control</b> .....  | <b>5</b>  |
| <b>Mode of Transmission</b> .....   | <b>6</b>  |
| <b>Pathology Specimens and Diagnosis of <i>Norovirus</i></b> .....  | <b>6</b>  |
| <i>Pathology Specimens</i> .....  | 6         |
| <i>Diagnosis</i> .....  | 6         |
| <b>Infection Control Precautions</b> .....  | <b>7</b>  |
| Contact and Droplet Precautions .....   | 7         |
| <i>Staff</i> .....  | 7         |
| <i>Parents and visitors</i> .....   | 7         |
| <b>Isolation and Placement of Patients with <i>Norovirus</i></b> .....  | <b>8</b>  |
| All inpatients .....  | 8         |
| Duration of Isolation Requirements for Specific Patient Groups .....  | 8         |
| <i>Bone marrow transplant (BMT) patients diagnosed with <i>Norovirus</i></i> .....                              | 8         |
| <i>General oncology or solid organ transplant patients diagnosed with <i>Norovirus</i></i> .....                | 8         |
| <i>Non-immunosuppressed patients who are diagnosed with <i>Norovirus</i></i> .....                              | 8         |
| <b>Parents and Carers of admitted patients</b> .....  | <b>9</b>  |
| <i>Parents and carers of children admitted with potential or proven infectious diarrhoea +/- vomiting</i> ..... | 9         |
| <i>Parents and carers who have gastroenteritis symptoms</i> .....   | 9         |
| <b>Hospital Volunteers</b> .....  | <b>9</b>  |
| <b>Patient Activity Outside Room</b> .....  | <b>10</b> |
| <b>Patient care equipment</b> .....   | <b>10</b> |
| <b>Room Cleaning</b> .....  | <b>11</b> |
| <i>Linen</i> .....  | 11        |
| <b>Eating Utensils</b> .....  | <b>11</b> |
| <b>Waste Management</b> .....   | <b>11</b> |
| <b>General Maintenance</b> .....  | <b>12</b> |
| <b>Discharge of Patient from Hospital</b> .....   | <b>12</b> |
| <b>Staff Management</b> .....   | <b>12</b> |
| Staff management during outbreak and ward closure situations. ....  | 12        |
| <b>Staff Education</b> .....  | <b>13</b> |
| Outbreak Management .....   | 13        |
| <b>Fact sheets</b> .....  | <b>13</b> |
| <i>Parent/Carer fact sheet</i> .....  | 13        |
| <i>Staff fact sheet</i> .....   | 13        |
| <b>References</b> .....   | <b>14</b> |

## Introduction

*Norovirus* is a leading cause of gastroenteritis worldwide and is recognised to be one of the most important causative agents responsible for gastroenteritis outbreaks in developed countries<sup>1</sup>. Low infectious dose, high infectivity, and short incubation period, extreme stability of the virus in the environment and frequent lack of prodromal symptoms associated with *Norovirus* infection render it an ideal agent for outbreak in confined environments such as hospitals, schools or institutions.

Outbreaks may occur at any time of the year, but most frequently during the winter months<sup>1</sup>.

## What is *Norovirus*?

Norovirus (genus *Norovirus*, family *Caliciviridae*) are a group of related, single-stranded RNA, non-enveloped viruses that cause acute gastroenteritis in humans. *Norovirus* was recently approved as the official genus name for the group of viruses provisionally described as "Norwalk-like viruses" (NLV). Currently, human noroviruses belong to one of three *Norovirus* genogroups (GI, GII, or GIV), each of which is further divided into >25 genetic clusters.

## Clinical Presentation

The incubation period for *Norovirus*-associated gastroenteritis in humans is usually 24 to 48 hours, but cases can occur within 12 hours of exposure<sup>2</sup>. *Norovirus* infection usually presents as acute-onset vomiting, watery non-bloody diarrhoea with abdominal cramps and nausea. Low-grade fever also occasionally occurs, and vomiting is more common in children. Dehydration is the most common complication, especially among the young and elderly, and may require medical attention. Symptoms usually last 24 to 60 hours. Recovery is usually complete and there is no evidence of any long-term sequelae. Studies have shown that as many as 30% of infections may be asymptomatic, although the role of such infections in *Norovirus* transmission is not well understood<sup>2</sup>.

## Command and Control

- Responsibility for implementation of this policy is the direct responsibility of appropriate clinical line managers caring for affected patients.
- The clinical line managers will consult with the infection control Infection Control Team regarding appropriate patient placement and infection control procedures.
- Where there is a dispute between clinical line managers and infection control / microbiology or if there is no policy on a particular issue or the policy needs updating then there needs to be further discussion between clinical line managers, infection control, microbiology and the Director of Clinical Operations to develop a consensus agreement based on best evidence. If a dispute arises about policy it is to be referred to the Chief Executive for resolution.
- *Norovirus* infection or colonisation is not mandated as a reportable infection to Public Health Units. Gastroenteritis amongst patients in the Hospital shall be notified to the Public Health Unit (PD2006\_014).
- A Reportable Incident Brief (RIB) will be sent to NSW Department of Health on any potential media interests or problems. This is currently the responsibility of the Executive Assistant to the CE.
- Microbiologist or Infection Control Practitioner will notify the Director of Clinical Operations of identification of any known *Norovirus* clusters. The Director of Clinical Operations will in turn notify the Chief Executive.
- A report on management of any new *Norovirus* cluster will be made to the next Infection Control Committee meeting. The Infection Control Committee minutes will be sent to the Health Care Quality Committee for information.
- Any ongoing outbreak of infections or colonisations not responding to appropriate infection control measures will be discussed with members of the Clinical Executive in collaboration with the appropriate clinical teams to discuss what further actions may be required.

## Mode of Transmission

*Norovirus* is transmitted primarily via the faecal-oral route, either by consumption of faecally contaminated food or water or by direct person-to-person spread. Environmental and fomite contamination may also act as a source of infection. Good evidence exists for transmission due to aerosolisation of vomitus that presumably results in droplets contaminating surfaces or entering the oral mucosa and being swallowed<sup>2</sup>.

*Norovirus* is highly contagious, and it is thought that an inoculum of as few as 10 viral particles may be sufficient to infect an individual<sup>2</sup>. Although pre-symptomatic viral shedding may occur, shedding usually begins with the onset of the symptoms and may continue for up to 2 - 3 weeks after recovery<sup>2</sup>. It is unclear to what extent viral shedding after clinical recovery signifies continued infectivity<sup>2</sup>.

## Pathology Specimens and Diagnosis of *Norovirus*

### *Pathology Specimens*

- Pathology personnel must comply with Contact and Droplet Precautions when entering and leaving the room.
- Stool samples are to be collected. **DO NOT** send vomitus to the laboratory as ELISA testing is not validated on this specimen type.
- Seal specimen receptacles correctly, label specimen accurately and complete all relevant details on the request form.
- Place specimen and pathology form into a plastic biohazard specimen bag for transport.
- Hand Hygiene with ABHR or antiseptic wash and water after sending specimen.

### *Diagnosis*

An ELISA assay on stool specimens is the test routinely used to diagnose *Norovirus* at our hospital. Reverse-transcriptase polymerase chain reaction (RT-PCR) is another way to test stool and emesis samples, as well as to detect the presence of *Norovirus* on environmental swabs. However RT-PCR is not performed at our hospital and can only be arranged after discussion with the Microbiologist / Virologist.

In addition to microbiological techniques, several epidemiological criteria have been proposed for use in determining whether an outbreak of gastroenteritis is of viral origin. Kaplan's criteria for this purpose are as follows:<sup>3</sup>

- a mean (or median) illness duration of 12 to 60 hours,
- a mean (or median) incubation period of 24 to 48 hours,
- more than 50% of people with vomiting and
- no bacterial agent identified on routine testing

## Infection Control Precautions

### Contact and Droplet Precautions

[Contact and droplet precautions](#) are to be used by **all staff** entering the child's room until the child has been asymptomatic for 72 hours.

Alcohol-based hand rubs (ABHR) may have slightly less activity against *Norovirus* than washing hands with water at a basin. High hand hygiene compliance is critical and during periods of sustained *Norovirus* transmission. A thorough hand wash with antiseptic wash and water needs to be done when dealing with patients that are symptomatic with *Norovirus*. The use of ABHR needs to remain routine for all healthcare staff to prevent transmission of other organisms.

#### **Staff**

- Hand hygiene
  - Use alcohol hand rub or antiseptic wash and water before entering the child's room.
  - Wash hands with antiseptic wash and water before leaving the room.
- Use alcohol hand rub antiseptic wash and water when outside the room before; leaving the ward or attending another patient. Personal Protective Equipment (PPE)
  - Wear gloves and disposable gowns when attending a patient who has diarrhoea and/or vomiting.
  - ADD surgical masks and eyewear if the patient is vomiting or is symptomatic with diarrhoea. Gloves must be worn when in contact with body fluids or blood of infected children as per standard precautions.
  - Gloves, gowns, surgical masks and eye protection are required when cleaning up vomit or faeces.
  - Remove PPE carefully without self-contamination on the inside of the room prior to leaving. Place disposable item in general waste bin provided in the room. Protective eye wear needs to be cleaned thoroughly with a 70% isopropyl wipe.

#### **Parents and visitors**

##### *Visitors should be limited to immediate family only*

- Hand hygiene
  - Use ABHR or antiseptic wash and water before entering the child's room.
  - Wash hands with antiseptic wash and water before leaving the room.
  - Use alcohol hand rub or antiseptic wash and water when outside the room before leaving the ward.
- Gloves, gowns and surgical masks are required when cleaning up vomit or faeces
- Toilets where body waste is being disposed should have the lid of the toilet closed before flushing to stop aerosols being generated.
- Parents and carers must follow hospital policy closely to avoid transmitting infection – see section below.

## Isolation and Placement of Patients with Norovirus

### All inpatients

- **Any child with *Norovirus*** should be nursed on the ward which is most appropriate for their medical needs.
- All children with *Norovirus* **MUST** be nursed in a single room or cohorted with other children with *Norovirus* in a dedicated room with en-suite toilet and bathroom facilities. **Only cohort after consultation with Infection Control.**
- Patients with *Norovirus* must not share a room or bathroom with patients who do not have *Norovirus*.
- The patient's room must have a staff hand wash basin.
- Adequate supplies of gowns, gloves, surgical masks and alcohol 'hand rub' are required outside the room.
- Patients who have been exposed but remained asymptomatic should only be isolated if they develop clinical symptoms.

**Note:** Notify Infection Prevention & Control team if there are any other patients, parents or carers with symptoms of gastroenteritis. Infection Control will notify the Public Health Unit if required.

### Duration of Isolation Requirements for Specific Patient Groups

#### ***Bone marrow transplant (BMT) patients diagnosed with Norovirus***

Because of possible prolonged excretion of *Norovirus*, BMT patients are considered infectious for the duration of that particular inpatient encounter at the Children's Hospital at Westmead. If they are to remain inpatients for an extended period of time, a clearance process for de-isolation may be considered. Before starting this process, the patient must be asymptomatic for at least 72 hours. Three stool samples collected one a week over three weeks, should be submitted to the laboratory for *Norovirus* testing. If all three samples test negative for *Norovirus*, the infection control status of the patient can be revised.

If they are discharged prior to this process, the stool screen does not need to be continued during the next encounter with the hospital if they have been asymptomatic for 72 hours.

- Must remain in isolation in a room with dedicated ensuite bathroom/toilet facilities.

#### ***General oncology or solid organ transplant patients diagnosed with Norovirus***

- Must be considered infectious while they have diarrhoea +/- vomiting.
- Once symptoms have resolved, can undergo a clearance process as described above for BMT patients.

#### ***Non-immunosuppressed patients who are diagnosed with Norovirus***

- When asymptomatic for 72 hours they can be taken out of isolation and maintained with standard precautions. **Follow-up stool testing is NOT required.**



## Parents and Carers of admitted patients

### ***Parents and carers of children admitted with potential or proven infectious diarrhoea +/- vomiting***

- Must not use shared facilities for food preparation in the ward or shared recreational areas in the ward or throughout the hospital even if they themselves are asymptomatic.
- Must not sleep in the parent hostel or parent rooms provided on the ward. If staying in the hospital with their child they must sleep in their child's room
- Must use the toilet and bathroom facilities in the child's isolation room.
- Must request nurse assistance to get food, beverages or feeding bottles from the Ward Kitchen for their child.
- If the parent needs to purchase meals themselves the parent can go to the providers in the hospital and either eat in an area isolated from other customers and patients - for example outdoor areas - or eat in their child's room. Any linen required by the patient or the parent must be provided by the nursing staff. Parents of symptomatic children are not to access the clean linen dispensary on the ward.

### ***Parents and carers who have gastroenteritis symptoms***

- Should be advised to stay home if possible.
- If they cannot stay home they must not use shared facilities for food preparation or shared recreational areas until asymptomatic for 72 hours.
- Must wash hands well with antiseptic hand-wash and water frequently, particularly after vomiting, after using the toilet, on leaving the patient's room, and before food or drink preparation.
- When they leave the child's room they must go straight home and not use shared Hospital facilities.
- Any linen required by the patient or the parent must be provided by the nursing staff. Parents of symptomatic children are not to access the clean linen dispensary on the ward.
- If there is a need to purchase meals, after the parents acute symptoms abate, the parent must liaise with the ward Nursing Unit Manager and After Hours Nurse Manager so that they can be assisted with this task while waiting for the 72 hours post resolution of symptoms to be attained.
- Also refer to the [Fact sheet for Parents and Carers on Norovirus Gastroenteritis](#).

## Hospital Volunteers

General visiting by hospital volunteers needs to be postponed until the patient or the symptomatic parent/carer has been symptom free for 72 hours.

There are some circumstances in which volunteer assistance is acceptable. In this case the volunteer needs to comply with the same requirements for hand hygiene and PPE usage as staff.

Ward Grandparent Volunteers can continue to work with their symptomatic child but need to comply with the same requirements for hand hygiene and PPE usage as parents.

Book Bunker lending should be postponed until the child or the symptomatic parent/carer has been symptom free for 72 hours.

Visitors organised by the Public Relations Department to the wards must not visit a symptomatic patient. This also must be postponed until the patient or the symptomatic parent/carer has been symptom free for 72 hours.

## Patient Activity Outside Room

- The child can use the outside areas in the hospital grounds.
- The child cannot visit the common food outlet areas.
- The child cannot visit the Starlight Room.
- The child cannot visit Ronald McDonald House.
- The child cannot attend the schoolroom.
- The child cannot visit other inpatients.
- Activities and school can be organised in the room.
- All other activities must be negotiated with Infection Control.

## Patient care equipment

- Must be dedicated for the **sole purpose** of the patient.
- The patient should have his / her own equipment such as stethoscopes, sphygmomanometers, thermometers and pans.
- This equipment should remain in the patient's room for the duration of the patients stay.
- Once the patient has left the room, all dedicated equipment must be wiped over with chlorine-based disinfectant (e.g. 1 dichloroisocyanurate tablet dissolved in 5L water for stainless steel surfaces, 1 tablet in 1L of water for other non-porous surfaces, and 5 tablets in 1L for porous surfaces). Other options include the autoclave or discarding the item.
- If the equipment has been in contact with faeces or vomit, it must be washed with a neutral detergent prior before cleaning with chlorine-based disinfectant as described above<sup>1, 4</sup>.
- For special equipment that cannot be cleaned according to the instructions above, it is important to minimise any possible exposure to *Norovirus* in the patient room. Other cleaning options can be discussed with the Infection Prevention and Control team.

## Room Cleaning

PPE, including gloves, gown, eye protection and a surgical mask, should be worn by people cleaning areas contaminated by faeces or vomit.

When cleaning the bathroom areas, special attention should be given to cleaning all potentially contaminated areas, including the toilet roll dispensers, toilet seats and lid, flushing mechanism, safety handles, shower chair, light switches, regardless of whether they are visibly soiled or not<sup>4</sup>.

Once the patient is discharged, the room should be cleaned, as per the Cleaning Services protocol.

### **Linen**

- All staff must perform hand hygiene immediately prior to accessing the ward's clean linen dispensary to prevent contaminating clean linen.
- If linen is removed from the clean linen dispensary it must not be replaced back onto the trolley, but be placed in to the used linen skip.
- PPE should be worn by staff when handling soiled linen from an infected patient, regardless of the child being in the bed or not.
- Used linen, whether visibly soiled or not, should not be shaken.
- Used linen should be bagged and tied at the point of generation. Care needs to be taken not to overfull the linen skip. It should not be filled more than  $\frac{3}{4}$  full so that it can be secured safely.
- The laundering of used linen should be consistent with Australian Standard AS 4146: Laundry Practice.

## Eating Utensils

Meal trays and eating utensils/plates and cups are to be collected from the room by staff with care. They can be placed in the Food Services trolley to be taken down to the Food Services department so they can be washed as per Food Services policy.

After carefully placing the used meal tray on the trolley staff need to be mindful to perform hand hygiene with antiseptic wash and water in case of viral contaminants on the tray.

## Waste Management

Toilets where body waste is being disposed should have the lid of the toilet closed before flushing to stop aerosols being generated.

General waste from a *Norovirus* patient's room is to be placed appropriately into the general waste receptacle. It is not to be over filled. When there is a requirement for a larger general waste bin to cope with the use of disposable gowns contact the cleaning services supervisor

so that an appropriate general waste bin can be obtained. After general working hours if the bin is more than  $\frac{3}{4}$  full contact the after-hours cleaning supervisor so that appropriate action can be taken.

## General Maintenance

Routine maintenance needs to be postponed until the patient has been symptom free for 72 hours.

Urgent maintenance can proceed with appropriate PPE wear and hand hygiene while the patient is in the acute stage of the illness.

Contact the Infection Prevention & Control team for advice if required.

## Discharge of Patient from Hospital

Discussion should take place before discharge to ensure the patient and family is fully informed about *Norovirus*. The patient should be requested to alert staff of *Norovirus* status if admitted to a health care facility in the following week. Children cannot be immediately discharged to Ronald MacDonald House, other housing arrangements must be organised until they are deemed non-infectious.

## Staff Management

Minimise as much as possible the circulation of staff between affected and unaffected areas<sup>1</sup>. Where possible, designated staff should care for affected patients.

Staff with gastrointestinal symptoms should leave work immediately and not return to work until 48 hours after their last episode of vomiting or diarrhoea<sup>4</sup>. Affected staff should seek medical advice immediately.

Food handlers should be excluded from food preparation until at least 72 hours after their symptoms have stopped<sup>4</sup>.

Recuperating staff may shed the virus for a number of weeks after their symptoms have disappeared, therefore the importance of hand washing and personal hygiene on returning to work should be reinforced<sup>4</sup>.

Non-essential staff should not be allowed to enter the patient care area of infected patients in order to prevent unnecessary exposure and to stop further spreading of the disease<sup>1</sup>.

### Staff management during outbreak and ward closure situations.

Non affected staff may leave the ward during their designated meal breaks. They need to ensure they hand wash with antiseptic wash and water immediately prior to leaving the ward, take care opening the main ward door, and use ABHR on the outside of the ward. They then may purchase meals and consume them in any recreational or dining area in the hospital.

For designated meal breaks taken on the ward, staff need to ensure they hand wash with antiseptic wash and water immediately prior to entering the area used as a staff meal/recreational room. ABHR should be used on the inside of the room prior to consumption of food and drink.

No food or drink other than water should be consumed in the ward area by staff including the 'nurses' station'.

Staff that have had cars valet parked earlier in the shift can leave the ward and go to the Security Department Office to get their keys during an evening shift; or prior to leaving at the end of the evening shift. However they must ensure they hand wash with antiseptic wash and water immediately prior to leaving the ward, take care opening the main ward door, and use ABHR on the outside of the ward.

## Staff Education

- Infection control will provide education on request.

Also refer to the [Norovirus Fact sheet for staff](#).

## Outbreak Management

If there appears to be two or more individuals in a ward area affected by symptoms indicative of Norovirus:

- Patients
- Parent/Carer and family/visitors
- Health Care Staff/Volunteers

The most senior staff member on the ward needs to refer to the [Gastro Cluster Checklist](#) which is located on the intranet in the Infection Prevention & Control department's page. The items in this check list needs to be actioned.

## Fact sheets

### *Parent/Carer fact sheet*

- [http://chw.schn.health.nsw.gov.au/ou/infection\\_control/resources/factsheets/parents/norovirus\\_gastroenteritis.pdf](http://chw.schn.health.nsw.gov.au/ou/infection_control/resources/factsheets/parents/norovirus_gastroenteritis.pdf)

### *Staff fact sheet*

- [http://chw.schn.health.nsw.gov.au/ou/infection\\_control/resources/factsheets/staff/norovirus.pdf](http://chw.schn.health.nsw.gov.au/ou/infection_control/resources/factsheets/staff/norovirus.pdf)

## References

1. Cheng, F.W.T., Leung, T.F., Lai, R.W.M., Chan, P.K.S., Hon, E.K.L. and NG, P.C. Rapid control of Norovirus gastroenteritis outbreak in an acute paediatric ward. *Acta Paediatrica*, 2006; 95:581-586
2. CDC – National Centre for Infectious Diseases – Respiratory and Enteric Viruses Branch. Norovirus Question and Answer sheet. <http://www.cdc.gov/ncidod/dvrd/revb/gastro/norovirus-qa.htm>
3. Turcios, RM.;Widdowson, M-A.;Sulka, A.; Glass, RI. Reassessment of Kaplan's criteria in identifying foodborne Norovirus outbreaks – United States, 1998–2000.
4. New South Wales Department of Health. 2007. Infection Control Policy PD2007 036: [http://www.health.nsw.gov.au/policies/pd/2007/pdf/PD2007\\_036.pdf](http://www.health.nsw.gov.au/policies/pd/2007/pdf/PD2007_036.pdf)
5. The Children's Hospital at Westmead Handbook: clinical practice guidelines for paediatrics / Henry Kilham and David Isaacs, editors. McGraw Hill, North Ryde, NSW, Australia
6. Australian Commission on Safety and Quality in Healthcare. Australian Guidelines for the Prevention and Control of Infections in Healthcare, 2010

### **Copyright notice and disclaimer:**

The use of this document outside Sydney Children's Hospitals Network (SCHN), or its reproduction in whole or in part, is subject to acknowledgement that it is the property of SCHN. SCHN has done everything practicable to make this document accurate, up-to-date and in accordance with accepted legislation and standards at the date of publication. SCHN is not responsible for consequences arising from the use of this document outside SCHN. A current version of this document is only available electronically from the Hospitals. If this document is printed, it is only valid to the date of printing.