

ACHONDROPLASIA: PRACTICE GUIDELINES FOR ALLIED HEALTH PROFESSIONALS

POLICY[®]

DOCUMENT SUMMARY/KEY POINTS

- This guideline has been written to assist the interdisciplinary team in providing care to infants and young children with Achondroplasia, it will also serve as a resource for collaborating community therapists
- Children with Achondroplasia have different patterns of growth and development to average statured people and this document outlines the normal progress expected for a child with Achondroplasia
- Children with Achondroplasia have unique risks of adverse musculoskeletal and neurological outcomes and need specific management to reduce these risks, which is outlined in this document.
- Children with rare skeletal dysplasia are also at risk of spinal deformity and subsequent neurological compromise, parts of this practice guideline may be recommended by their medical specialist to assist with their care on a case by case basis.

This document reflects what is currently regarded as safe practice. However, as in any clinical situation, there may be factors which cannot be covered by a single set of guidelines. This document does not replace the need for the application of clinical judgement to each individual presentation.

Approved by:	SCHN Policy, Procedure and Guideline Committee	
Date Effective:	1 st December 2016	Review Period: 3 years
Team Leader:	Staff Specialist	Area/Dept: Kids Rehab

CHANGE SUMMARY

- N/A – new document.

READ ACKNOWLEDGEMENT

- Any Health professional providing clinical care to babies and infants with Achondroplasia needs to read and be aware of this whole document
- This document is NOT APPROPRIATE for parents and families – please see separate factsheet

This document reflects what is currently regarded as safe practice. However, as in any clinical situation, there may be factors which cannot be covered by a single set of guidelines. This document does not replace the need for the application of clinical judgement to each individual presentation.

Approved by:	SCHN Policy, Procedure and Guideline Committee	
Date Effective:	1 st December 2016	Review Period: 3 years
Team Leader:	Staff Specialist	Area/Dept: Kids Rehab

TABLE OF CONTENTS

1	Introduction and Background	4
2	What is Achondroplasia?	4
2.1	What musculoskeletal and medical issues need to be considered when treating a child with achondroplasia?.....	4
2.1.1	<i>Spinal Canal Stenosis</i>	4
2.1.2	<i>Thoracolumbar Kyphus</i>	5
2.1.3	<i>Elbow Flexion</i>	5
2.1.4	<i>Lower limb deformity</i>	5
2.1.5	<i>Joint pain and fatigue</i>	6
2.1.6	<i>Hearing</i>	6
2.1.7	<i>Sleep Apnoea</i>	6
2.1.8	<i>Hydrocephalus</i>	6
2.2	Gross motor development.....	6
2.3	Fine motor development.....	8
2.4	Communication skills.....	8
2.5	Feeding behaviour.....	8
3	Musculoskeletal Management	9
3.1	Protection of the Newborn Cervical Spine.....	9
3.2	Prevention of Fixed Thoracolumbar Kyphus.....	9
3.3	Management of Elbow & Forearm Deformity.....	10
4.1	Management.....	11
5	Childcare Preparation	12
6	References	12

1 Introduction and Background

These guidelines have been developed to assist health professionals when treating babies and children with Achondroplasia. The CMSK rehab team consists of; a rehabilitation physician, physiotherapist and occupational therapist, and is part of the specialist skeletal dysplasia clinic based at the Children's Hospital at Westmead. This group are also treated and provided with therapy at SCH, where appropriate (eg hydrotherapy).

The team aims to review patients with achondroplasia every 4 months in the first year of life, then every 6 months until 3 years of age, then yearly following that until starting school.

The clinic works in collaboration with local medical and therapy teams to deliver best outcomes for the children, with as much care as close to home as possible.

Please do not hesitate to contact any of the team with any questions, queries or feedback.

Contact details:	
Physiotherapist	9845 2132
Occupational Therapist	9845 3369
Rehabilitation Physician	9845 2132

2 What is Achondroplasia?

Achondroplasia is a congenital skeletal dysplasia affecting approximately 1 in every 25,000 births, resulting in disproportionate short stature. Children with achondroplasia have normal sitting height (trunk length), a large head and shortened long bones, particularly the humerus and femur. Other skeletal anomalies include: restricted elbow extension, supination, short hands and fingers, short broad and flat feet. Hypermobility and ligament laxity of the fingers, wrists, feet, knees and spine are usually present.

Average adult heights are approximately 130cm (males) and 126cm (females). Intelligence is generally normal, and all children will be expected to participate in normal education pathways.

2.1 What musculoskeletal and medical issues need to be considered when treating a child with achondroplasia?

2.1.1 Spinal Canal Stenosis

Children with achondroplasia have a small posterior cranial fossa and narrowing of the spinal canal (1/2 – 1/3 of the size of their non-affected peers) as a result of shortened pedicles and thickened laminae. The brainstem and spinal cord are normal size, so the fit inside the canal is quite tight, particularly at the foramen magnum. Pressure on the brainstem or spinal cord may cause symptoms of spinal canal stenosis or high cervical myelopathy.

Hip flexion contractures or poor core stability result in a hyperlordotic lumbar posture which further reduces the space in the spinal canal for the spinal cord, and may worsen spinal canal stenosis. Increased weight for height will also increase the lumbar lordosis and place greater stress on the already narrowed spinal canal.

Symptoms may include:

High stenosis (brainstem or high cervical level) – central sleep apnoea, oromotor dysfunction including difficulties swallowing and articulating, central sleep apnoea, hyperreflexia, hypertonia or clonus

Low stenosis (thoracic or lumbar levels) – pain, altered sensation, weakness or intermittent claudication in lower limbs, bladder and bowel dysfunction, hyporeflexia

2.1.2 Thoracolumbar Kyphus

Anterior wedging of thoracolumbar vertebrae can develop in children with achondroplasia due to the bone dysplasia and ligamentous laxity of their spine. This allows a thoracolumbar kyphus to develop in the presence of muscle weakness (refer to fig 1). Ongoing muscle weakness may result in this kyphus becoming fixed, further exacerbating the spinal canal stenosis as the cord is stretched over the longer kyphotic length. Wedging is usually preventable.



Fig 1. Thoracolumbar Kyphus
<http://www.nemours.org/service/medical/orthopedics/dysplasia/achondroplasia/characteristic.html>

2.1.3 Elbow Flexion

The elbow positioning is bony. No attempt should be made to increase range by splinting.

2.1.4 Lower limb deformity

Children with achondroplasia commonly develop 'genu varum' (bow legs) as a result of their ligamentous laxity around the knee joint. This is normal and unless severe, will almost entirely resolve over time with increases in strength (refer to Fig 2).

Some children may develop tibial bowing, possibly including a leg length discrepancy, and may require surgical correction of the deformity. Surgery is rarely indicated in the first 5 years of life. Specialist orthopaedic consultation is available through the CMSK clinic.

Fig 2: Genu Varum – at rest and after asking the child to “stand up straight” illustrating the contribution of ligamentous laxity



2.1.5 Joint pain and fatigue

Joint pain and fatigue are common in the hypermobile joints of children with achondroplasia, particularly in the knees during walking and sporting activities, and in the hands during writing and fine motor tasks. Pain and fatigue problems are sometimes present in toddler years, although they tend to become more of an issue with the increasing school demands over time.

2.1.6 Hearing

The Eustachian tube is shorter and more horizontal, within the short skull base. Children with achondroplasia have a tendency toward middle ear infections in the first five to six years of life. Any infections must be recognized and treated promptly, and hearing should be checked on a regular basis to minimize any hearing loss. Children with an abnormal newborn hearing test should be referred under those guidelines and all children should have a repeat hearing test at 1 year old.

2.1.7 Sleep Apnoea

Children with achondroplasia are at an increased risk of obstructive sleep apnoea due to their small airways, and central sleep apnoea due to possible spinal canal stenosis and brainstem compression. Early referral (infant) is useful. Sleep studies should be performed regularly to monitor for this and appropriate weight for height should be maintained. Children with untreated sleep apnoea are likely to be easily fatigued and slower in their early development.

2.1.8 Hydrocephalus

As a baby with achondroplasia has a large head, hydrocephalus is often suspected. Medical monitoring of head growth on achondroplasia specific growth charts, and baseline head ultrasound should be performed. Surgical treatment is rarely necessary but may be indicated if the child has significant abnormal neurological symptoms.

2.2 Gross motor development

Babies with achondroplasia are likely to have a delay in gross motor skill acquisition compared to other children. Balancing a large head on a small hypermobile neck requires extra back and neck extensor strength, often resulting in a delay in obtaining head control. Current treatment aimed at preventing spinal deformity specifically delays independent and upright sitting, further contributing to gross motor delay. Children with achondroplasia also use unconventional methods to mobilize independently when young, often due to their shortened limbs, and do not follow “normal” developmental progress. For example, children with achondroplasia almost never play with their feet, as they are unable to reach them with their shortened limbs.

Fig 3



Snow-ploughing



Reverse snow-ploughing



Bear walking



Push to sit through prone



Move to sit through side-lying

Recent research on Australian children with achondroplasia, combined with our clinical experience leads us to consider the following progress to be within normal limits for these children. However infants with motor delay (compared to normal peers) must be screened at regular (4-6 monthly) intervals by their medical specialist to ensure the delay is not due to emerging neurosurgical issues:

- Able to lift head up in prone by 6 months
- Able to roll by 7 months
- Walk with hands held by 21 months
- Stand independently by 24 months
- Walk independently by 26 months

Children with achondroplasia also have very different movement patterns. When moving independently on the floor, traditional 4-point crawling is uncommon. Commando crawling is the most common form of movement for these children, although snow ploughing, reverse snow ploughing and bear walking are also common. All of these movements are normal and safe for children to perform (refer to Fig 3 above).

Transitional movements between positions often use very different movement patterns also. When moving from lying into sitting, most children with achondroplasia will lie prone, abduct their legs to almost 180°, and then push their trunks up into sitting. When pulling to stand, many children with achondroplasia will come up through half kneeling, with one leg abducted, and weight bear through a hyper-extended knee to reach standing. When moving from standing to sitting, many children will either just drop backwards or abduct their hips to get to the floor. All of these movement patterns are normal and safe for children with achondroplasia.

2.3 Fine motor development

The acquisition of fine motor skills in babies with achondroplasia is similar to that of other children. However, babies and toddlers with achondroplasia often have difficulties in grasping large objects due to their typically short fingers. Limited elbow extension and supination, combined with hyper-mobile wrist and finger joints can also create difficulties with some fine motor skills.

2.4 Communication skills

Babies and toddlers with achondroplasia may have a speech delay and difficulties with their expressive communication. There are many possible causes for this delay, including hearing problems and restricted floor play and delayed seating (for back care). Speech should be developing within normal limits, and a specialist opinion is needed if this is not the case for an individual child.

Please note:

13% of achondroplasia toddlers are late talkers (Less than fifty words and no phrases by 24 months). However, we hope that with better early intervention, this will improve.

2.5 Feeding behaviour

Children with achondroplasia tend to also have a delay in their feeding habits when compared to other children. Respiratory difficulties, relatively large tongue and small airways can create difficulties and noisy breathing when bottle feeding and children with achondroplasia will tend to have smaller volumes of feeds. However, it must always be remembered that poor oromotor dysfunction can also be a sign of high spinal cord compression including difficulties with coughing and gagging, spillage from the mouth and difficulty with new textures.

The following developmental progress is currently considered to be within normal limits for these children:

- Manage smooth solids at 4-7 months
- Manage mashed solids at 6-12 months

- Perform finger feeding 8-18 months
- Drink from a cup 8-22 months
- Self feed with a spoon 15-27 months

3 Musculoskeletal Management

Provide parents and carers with the [‘positioning and handling of babies with Achondroplasia’](#) handout developed by the Complex Musculoskeletal Therapists. This provides illustrations of correct positioning and handling and outlines precautions.

3.1 Protection of the Newborn Cervical Spine

Due to the large head size, head control usually takes a longer time to develop in babies with Achondroplasia. Protection of the hypermobile neck is essential to minimise the risk of high cervical myelopathy from spinal canal stenosis.

Recommendations and Precautions:

Educate all carers on providing appropriate support to the baby’s head and neck during handling, to protect the cervical spine. Avoid extreme flexion and extension of the cervical spine to minimise the risk of high cervical myelopathy from spinal canal stenosis.

Do not lift the baby up by the arms or pull-to-sit, with an unsupported head.

Carrying pouches and mechanical swings are not recommended due to the uncontrolled head movement and risk of central and obstructive apnoea.

3.2 Prevention of Fixed Thoracolumbar Kyphus

Although a fixed thoracolumbar kyphus due to anterior wedging of vertebrae was once extremely common, experience in our clinic and overseas indicates that by delaying upright sitting and promoting trunk extensor muscle strength, most cases of fixed deformity may be prevented. 90% of infants with Achondroplasia will have a flexible kyphosis in the first year of life, most resolve with this programme. In our centre about 5% of the group require thoracolumbar bracing compared to 30% in centres overseas who do not follow this protocol.

Recommendations and Precautions:

- Independent sitting and standing should not be encouraged until the child has sufficient back strength to actively maintain an extended upright posture, and is able to attain sitting and standing independently.
- Development of trunk extensor muscle strength should be actively encouraged with prone activities (tummy time). Wedges, therapy balls, rolled up towels etc. may be used.
- When babies are awake they should be positioned in prone, side lying or supine (prone is the preferred position to develop extensor muscle strength).
- Firm thoracolumbar support should be provided in reclined sitting and during handling until the child is able to sit independently. Flexed postures in seating should be avoided and a normal lumbar lordosis and spinal curve should try to be maintained.

- Babies should not be held or carried for more than a few minutes in a “curled-up” position. Baby slings and pouches are not recommended due to insufficient support of the thoracolumbar spine and risk of apnoea.
- “Burping” against the carers shoulder is a good position as the baby’s back is easy to support properly, especially if the carer is encouraged to lean back against pillows/cushions. Rubbing rather than patting is preferred when burping baby. Time in upright position should be limited.
- Sitting in a reclined seat is usually commenced around 4 months of age. This should be restricted to short periods of time and only used when the child is awake and alert. An infant seat with a firm back support such as the Frazer Chair is recommended. Initially the chair should be reclined back to 30 degrees from the ground. This angle can be gradually increased as strength and age allows.
- Car capsules, baby car seats and strollers should all have a firm back and be reclined as much as possible initially. Babies are best managed in a bassinet attachment for their prams. The seat or stroller can be placed in a more upright position over time as the baby gains better strength and head control.
- Seats to assist with bathing should all have a firm back and be reclined 30 degrees from the ground initially.
- Seats to assist with feeding eg high chair, should all have a firm back and be reclined to 60 degrees from the horizontal during feeding, until the child is able to sit independently with good trunk control.
- Aquatic physiotherapy activities may also help build muscle strength from approximately 6 months onwards (depending on head control and respiratory status).
- Baby bouncers, baby rockers, Jolly Jumpers, baby walkers and baby swings are not recommended due to extra force travelling through the unsupported spine.



Seating in a Fraser chair: foam is used to help position the Achondroplasia child safely, & to support back & neck.

3.3 Management of Elbow & Forearm Deformity

Limitations in elbow extension and supination result from bony changes (possible posterior humeral bowing and radial head dislocations). Most children with achondroplasia will continue to have functional range.

Recommendations:

- Monitor elbow joint range.
- Do not cast or attempt stretching to change the deformity. Reduced range is due to bony changes and will not be improved with casting or stretches.

4 Promotion of Early Development

As outlined above, children with achondroplasia do not follow “normal” developmental sequences and patterns. Standardised developmental assessments should not be used in this patient group. Wherever possible, parents should be provided with reassurance that their child’s movement patterns are safe and normal for Achondroplasia. Penny Ireland’s Developmental Screening Form is useful to monitor development compared to the Achondroplasia population and to provide reassurance to parents⁴. Education and support in promoting early language skills, along with encouraging opportunities for fine motor skill practise, while still maintaining positioning and handling to ensure best possible spine care, is important for families. The accompanying parent factsheet has language stimulation suggestions for families.

4.1 Management

In the first couple of years, prevention of musculoskeletal deformity needs to take priority over “normal” gross motor development. However, fine motor, speech and language development and stimulation should continue as per “normal”.

Provide education and support to parents on the need for musculoskeletal prevention to be paramount at this stage. Explain that their child will walk and sit like other children when they are ready, which will be a little later than average statured children. Many parents find this very difficult to accept, particularly when attending playgroups or around other children of a similar age.

Do not push or encourage the acquisition of sitting, standing or weight bearing positions until trunk strength is sufficient to prevent kyphotic posture.

Encourage speech and language development – encourage parents to talk, sing, laugh and interact face to face with their baby. Provide a stimulating physical environment, but allow children to develop at their own pace.

Assist the family in finding appropriate small, lighter toys, for their baby’s small hands to manage.

Provide opportunities for fine motor developmental play eg sitting reclined in a Fraser chair with a table around to allow for fine motor play a child would normally do in an independent sitting position. Crayons or pencils for drawing, or blocks for play should be easily held in their small hypermobile hands.

Encourage floor play, especially prone time, from an early age. Supine, side lying and prone should all be used for play, along with reclined seating.

Reassure parents when their child performs unconventional means of transitioning and moving.

Once the child has sufficient head control, aquatic physiotherapy may be commenced to provide another environment for strengthening and promoting motor skills. Supported sitting and standing can be trained in neck deep water if spinal posture is appropriately maintained.

Always encourage age-appropriate play and language. Remember that the achondroplasia child often looks much smaller and younger than they really are. Assist families not to 'baby' their child, but provide age appropriate activities for them. Provide advice on how to play and communicate with their child.

5 Childcare Preparation

Due to the risk of spinal complications childcare staff need to be educated correct positioning and handling and restrictions of the infant with Achondroplasia. Please contact the Complex Musculoskeletal Team at The Children's Hospital at Westmead for advice.

6 References

1. Brinkman G. Schlitt H. Zorowka P. Spranger J. Cognitive skills in achondroplasia. American Journal of Medical Genetics. 1993; 47: 800-804.
2. Fowler ES. Gliniski LP. Reiser CA. Horton VK. Pauli RM. (1997). Biophysical bases for delayed and aberrant motor development in young children with achondroplasia. Developmental and Behavioural Paediatrics. 1997; 18(3): 143-150.
3. Hunter AGW. Bankier A. Rogers JG. Silence D. Scott CL. Medical complications of achondroplasia: A multicentre patient review. Journal of Medical Genetics. 1998; 35: 705-712.
4. Ireland P. O'Donaghey S. Development of Australian children with achondroplasia. Journal of Developmental and Behavioural Paediatrics. 2007; 31(1): 41-7.
5. Thompson NM. Hecht JT. Bohan TP. Kramer LA. Davidson K. Brandt ME. Fletcher JM. Neuroanatomic and neurophysiological outcome in school-aged children with achondroplasia. American Journal of Medical genetics. 1999; 88: 145-153.
6. Todorov AB. Scott CI. Warren AE. Leeper JD. Developmental screening tests in achondroplastic children. American Journal of Medical Genetics. 1981; 9: 19-23.
7. Trotter TL. Hall JG & American Academy of Pediatrics Committee on Genetics. Health supervision for children with achondroplasia. Pediatrics. 2005; 116: 771-783.

Copyright notice and disclaimer:

The use of this document outside Sydney Children's Hospitals Network (SCHN), or its reproduction in whole or in part, is subject to acknowledgement that it is the property of SCHN. SCHN has done everything practicable to make this document accurate, up-to-date and in accordance with accepted legislation and standards at the date of publication. SCHN is not responsible for consequences arising from the use of this document outside SCHN. A current version of this document is only available electronically from the Hospitals. If this document is printed, it is only valid to the date of printing.