

## Lung Ultrasound in Children With COVID-19

Marco Denina, Carlo Scolfaro, Erika Silvestro, Giulia Pruccoli, Federica Mignone, Marisa Zoppo, Ugo Ramenghi, Silvia Garazzino

**DOI:** 10.1542/peds.2020-1157

**Journal:** *Pediatrics*

**Article Type:** Research Brief

**Citation:** Denina M, Scolfaro C, Silvestro E, et al. Lung ultrasound in children with COVID-19. *Pediatrics*. 2020; doi: 10.1542/peds.2020-1157

This is a pre-publication version of an article that has undergone peer review and been accepted for publication but is not the final version of record. This paper may be cited using the DOI and date of access. This paper may contain information that has errors in facts, figures, and statements, and will be corrected in the final published version. The journal is providing an early version of this article to expedite access to this information. The American Academy of Pediatrics, the editors, and authors are not responsible for inaccurate information and data described in this version.

## Lung Ultrasound in Children With COVID-19

*Marco Denina<sup>1</sup>; Carlo Scolfaro<sup>1</sup>; Erika Silvestro<sup>1</sup>; Giulia Pruccoli<sup>1</sup>; Federica Mignone<sup>1</sup>; Marisa Zoppo<sup>1</sup>; Ugo Ramenghi<sup>1</sup>; Silvia Garazzino<sup>1</sup>*

1. Department of Pediatrics, Infectious Diseases Unit, University of Turin, Regina Margherita Children's Hospital, Turin, Italy

Correspondence: Marco Denina, MD, Department of Pediatrics, Infectious Diseases Unit, University of Turin, Regina Margherita Children's Hospital, Piazza Polonia 94, 10126 Turin, Italy. Tel: 39 011 3135396 . Fax: 39 011 3135473 . E-mail: [marco.denina@gmail.com](mailto:marco.denina@gmail.com)

Financial Disclosure: The authors have no financial relationships relevant to this article to disclose.

Funding Source: No external funding for this manuscript.

Potential Conflicts of Interest: The authors have no conflicts of interest relevant to this article to disclose

Abbreviations:

LUS: Lung ultrasound

COVID-19: Coronavirus Disease 2019

Dr Denina conceptualized and designed the study, performed lung ultrasound examinations, drafted the initial manuscript, and revised the manuscript.

Drs Scolfaro, Pruccoli, Zoppo and Mignone, collected data, carried out the initial analyses, and reviewed and revised the manuscript.

Prof Ramenghi, Drs Garazzino and Silvestro supervised data collection, and critically reviewed the manuscript.

All authors approved the final manuscript as submitted and agree to be accountable for all aspects of the work.

## Introduction

The novel Coronavirus Disease 2019 (COVID-19) is rapidly spreading all around the world, affecting both adults and children. Recently, the clinical and radiographic characteristics of infected children have been described<sup>1</sup>. Even though lung ultrasound (LUS) is recognized as a valid imaging technique for the diagnosis and follow-up of pneumonia in children<sup>2</sup>, no data are currently available about LUS use in COVID-19 affected children. Considering the well-known advantages of point-of-care ultrasound<sup>3</sup>, including the possibility of reducing the patient's movement across the hospital department, we investigated LUS findings in children infected with Sars-Cov2.

## Methods

We performed an observational study at Regina Margherita Children Hospital (Turin, Italy). Children and adolescents (0-17 years) admitted to the pediatric infectious diseases department for documented COVID-19 were analyzed. Epidemiological, clinical, and laboratory data were collected, and we categorized patients according to the clinical definitions recently provided by Qiu et al<sup>1</sup>. LUS was performed during the routine medical examination, with a linear array transducer at 7.5 to 13 MHz (MyLab™Seven; Esaote) applying the Copetti protocol<sup>4</sup>.

## Results

Between March 18 and March 26, 2020, eight children affected by COVID-19 were admitted to our department. Epidemiological and clinical features are provided in table 1. Two patients (25%) were classified as severe clinical types, two (25%) as moderate, and four (50%) as mild. Two (25%) children needed non-invasive oxygen administration for 2 and 4 days, respectively. None of them required mechanical ventilation. We performed LUS in all eight patients, paying

particular attention to signs of viral pneumonia as small subpleural consolidations and/or individual B-lines or confluent B-lines (echogenic vertical lines arising from the pleural line and moving in concert with a sliding lung, expression of an interstitial syndrome)<sup>5</sup>. LUS documented subpleural consolidations in 2 children and confluent B-lines in 5. In 7/8 cases, we found a concordance with the radiologic findings (figure 1), whilst in the remaining patient, an interstitial B-lines pattern was observed despite a normal chest radiography. One patient with severe clinical type was repeatedly examined with LUS on alternate days, and we noted a B-lines bilateral pattern reduction a day in advance before clinical and radiographic improvement.

In the seven children with pathologic ultrasound imaging at baseline, LUS was repeated before discharge, showing improvement or resolution of consolidations (2/2) and interstitial patterns (5/5), consistent with the concomitant radiologic findings.

## Conclusion

Investigators have shown that in adults with COVID-19 pneumonia, bedside US correlates with CT findings<sup>6</sup>. Our study represents a preliminary report of lung ultrasound characteristics in children affected by COVID-19. While the number of patients analyzed is small, the high concordance between radiologic and LUS findings suggests that ultrasound may be a reasonable method to detect lung abnormalities in children with COVID-19. LUS may be useful in the management of children with COVID-19 for several reasons. First, it may reduce the number of radiologic examinations, lowering the radiation exposure of the patients. Secondly, being performed at the bedside, it allows to reduce the patient's movement within the hospital, thus lowering the number of health-care workers and medical devices exposed to Sars-Cov2. Moreover, the recent evolution in the ultrasound field allows the use of wireless devices, which, when available, are probably the most appropriate ultrasound equipment in patients with

confirmed or suspected COVID-19. Both the wireless probe and the tablets are easily wrapped in disposable plastic covers, allowing simple sterilization procedures and reducing the risk of contamination, as recently suggested by Buonsenso et al<sup>7</sup>.

In conclusion, although further studies are needed to better understand and characterize LUS findings in this novel disease in children, we propose LUS routinely protocol examinations as a useful tool in the diagnostic and clinical management of mild/severe Covid19 in children.


**Acknowledgments:** to Drs. AG. Del Monaco and E. Garrone, for their constant ultrasound teaching activity at pediatrics resident.

## References

1. Qiu H, Wu J, Hong L, Luo Y, Song Q, Chen D. Clinical and epidemiological features of 36 children with coronavirus disease 2019 (COVID-19) in Zhejiang, China: an observational cohort study. *Lancet Infect Dis*. March 2020. doi:10.1016/S1473-3099(20)30198-5
2. Pereda MA, Chavez MA, Hooper-Miele CC, et al. Lung ultrasound for the diagnosis of pneumonia in children: a meta-analysis. *Pediatrics*. 2015;135(4):714-722. doi:10.1542/peds.2014-2833
3. Lichtenstein DA, Mezière GA. Relevance of Lung Ultrasound in the Diagnosis of Acute Respiratory Failure\*: The BLUE Protocol. *Chest*. 2008;134(1):117-125. doi:10.1378/chest.07-2800
4. Copetti R, Cattarossi L. Ultrasound diagnosis of pneumonia in children. *Radiol Med*. 2008;113(2):190-198. doi:10.1007/s11547-008-0247-8
5. Tsung JW, Kessler DO, Shah VP. Prospective application of clinician-performed lung ultrasonography during the 2009 H1N1 influenza A pandemic: distinguishing viral from bacterial pneumonia. *Critical Ultrasound Journal*. 2012;4(1):16. doi:10.1186/2036-7902-4-16
6. Poggiali E, Dacrema A, Bastoni D, et al. Can Lung US Help Critical Care Clinicians in the Early Diagnosis of Novel Coronavirus (COVID-19) Pneumonia? *Radiology*. March 2020:200847. doi:10.1148/radiol.2020200847
7. Buonsenso D, Pata D, Chiaretti A. COVID-19 outbreak: less stethoscope, more ultrasound. *The Lancet Respiratory Medicine*. March 2020. doi:10.1016/S2213-2600(20)30120-X

**Table 1:** Epidemiological and clinical features of COVID-19 pediatric patients stratified by clinical types.

	Total (n=8)	Severe cases (n=2)	Mild cases (n=2)	Moderate cases (n=4)
<b>Epidemiological data</b>				
Female patients	3 (37.5%)	1 (50%)	1 (50%)	1 (25%)
Male patients	5 (62.5%)	1 (50%)	1 (50%)	3 (75%)
Age, years (SD, range)	4.2 (0.2-10)	3.9 (3.9 – 4)	5.1 (0.2-10)	3.8 (0.5 – 7.2)
Family members with COVID19	8 (100%)	2 (100%)	2 (100%)	4 (100%)
<b>Symptoms</b>				
Fever (body temperature >37.5°C)	6 (75 %)	2 (100%)	2 (100%)	2 (50%)
Dry cough	5 (62.5%)	2 (100%)	2 (100%)	1 (25%)
Dyspnea / tachypnea	3 (37.5%)	2 (100%)	1 (50%)	0
Pharyngeal congestion or sore throat	3 (37.5%)	1 (50%)	1 (50%)	1 (25%)
Vomiting or diarrhea	3 (37.5%)	0	2 (100%)	1 (25%)
<b>Hypoxia</b>	2 (25%)	2 (100%)	0	0
<b>Radiography</b>				
Pulmonary consolidation	2 (25%)	1 (50%)	1 (50%)	
Ground-glass opacities	4 (50%)	2 (100%)	1 (50%)	1 (25%)
<b>Lung Ultrasound</b>				
Pulmonary consolidation	2 (25%)	1 (50%)	1 (50%)	0
Interstitial B-lines pattern	5 (62.5%)	2 (100%)	2 (100%)	1 (25%)



**Figure 1:** In image C chest radiography with left upper lobe consolidation and right central ground-glass opacities; in images A and B multiple confluent B-lines (thin arrows) and in images D and E a lung consolidation (thick arrows) at lung ultrasound.

## Lung Ultrasound in Children With COVID-19

Marco Denina, Carlo Scolfaro, Erika Silvestro, Giulia Pruccoli, Federica Mignone, Marisa Zoppo, Ugo Ramenghi and Silvia Garazzino  
*Pediatrics* originally published online April 21, 2020;

### Updated Information & Services

including high resolution figures, can be found at:  
<http://pediatrics.aappublications.org/content/early/2020/04/17/peds.2020-1157.citation>

### Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:  
<http://www.aappublications.org/site/misc/Permissions.xhtml>

### Reprints

Information about ordering reprints can be found online:  
<http://www.aappublications.org/site/misc/reprints.xhtml>

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



# PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

## **Lung Ultrasound in Children With COVID-19**

Marco Denina, Carlo Scolfaro, Erika Silvestro, Giulia Pruccoli, Federica Mignone, Marisa Zoppo, Ugo Ramenghi and Silvia Garazzino  
*Pediatrics* originally published online April 21, 2020;

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/early/2020/04/17/peds.2020-1157.citation>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2020 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

